

Adenanthera pavonina

RED SANDALWOOD

Fabaceae

Common Synonyms: *none*

FLEPPC Category: 2

FDACS Listed Noxious Weed: No


IFAS Assessment

North	OK
Central	OK
South	CAUTION

USDA Hardiness Zone: 10B - 11

Growth Habit: Tree

Origin: Tropical Asia, S. China, India,
N. Australia, and Soloman Islands



Mauricio Mercadante

Description: Deciduous tree to 20 m tall with a buttressed trunk and flaky bark. Alternate, stalked, leaves, twice compound to 40 cm long, glabrous to glaucous beneath, dark green above. Small pale yellow to orange flowers (not pea-shaped) on elongated stalks arising from leaf axils, fragrant. Fruit a long narrow pod, maturing black and twisted with shiny red seeds.

Note: Similar to *Albizia lebbek*, bark of *Adenanthera pavonina* reddish

Habitat: Prairie hammock, rockland hammock, disturbed sites

Comments: Potential problem species in tropical areas, possible EDRR candidate if not well established. Seeds persist in soil for up to 5 years. Seeds dispersed by birds.

Florida Introduction Date: pre-1930

Control Methods:

Mechanical: hand-pull seedlings (IFAS).

Chemical: Basal bark (10% triclopyr ester, IFAS), cut stump (50% triclopyr amine or 10% triclopyr ester, IFAS).

Useful Resources:

Dave's Garden. 2013. PlantFiles: Red sandalwood, Coral tree, *Adenanthera pavonina*. <http://davesgarden.com/guides/pf/go/51496/>. Accessed on December 2, 2013.

Langeland, K.A., H.M. Cherry, C.M. McCormick, K.C. Burks. 2008. Identification and Biology of Non-Native Plants in Florida's Natural Areas-Second Edition. IFAS Publication SP 257. University of Florida, Gainesville, Florida.