Dioscorea bulbifera

AIR-POTATO

Dioscoreaceae

Common Synonyms: none

FLEPPC Category: 1


FDACS Listed Noxious Weed: Yes

IFAS Assessment

North PROHIBITED
Central PROHIBITED
South PROHIBITED

USDA Hardiness Zone: 9a-11

Growth Habit: Vine **Origin:** Tropical Asia





Description: Twining, herbaceous vine, with small or absent underground tubers. Stems to 20 m or more. Internodes, round, not winged. Bulbils roundish to 12 cm, surfaces smooth and formed in leaf axils. Leaves with long stalks, alternate, to 20 cm, broadly heart-shaped. Flowers rare, small fragrant, dioecious, in panicles or spikes to 11 cm long. Fruit a capsule.

Note: Similar to D. alata which has opposite leaves and native D. floridana, which never forms bulbils and has smaller leaves.

Habitat: hammocks, hardwood forests, disturbed areas

Florida Introduction Date: 1905

Control Methods: Mechanical: Cut vines that are high in trees; cut bulbils and remove from site. Dig up underground tubers if possible..

Chemical: Foliar (1-2% glyphosate [IFAS]), cut stems (10% triclopyr ester [IFAS] or undiluted triclopyr amine [UGA])..

Useful Resources:

Langeland, K.A., H.M. Cherry, C.M. McCormick, K.C. Burks. 2008. Identification and Biology of Non-Native Plants in Florida's Natural Areas-Second Edition. IFAS Publication SP 257. University of Florida, Gainesville, Florida.

Langeland, K.A., J.A. Ferrell, B. Sellers, G.E. MacDonald, and R.K. Stocker. 2011. Integrated management of non-native plants in natural areas of Florida. EDIS publication SP 242. University of Florida, Gainesville, Florida.

Wunderlin, R. P., and B. F. Hansen. 2008. Atlas of Florida Vascular Plants (http://florida.plantatlas.usf.edu/).[S. M. Landry and K. N. Campbell (application development), Florida Center for Community Design and Research.] Institute for Systematic Botany, University of South Florida, Tampa.

Comments: