

Paederia foetida

SKUNKVINE

Rubiaceae

Common Synonyms: none

FLEPPC Category: 1

FDACS Listed Noxious Weed: Yes


IFAS Assessment

North	PROHIBITED
Central	PROHIBITED
South	PROHIBITED

USDA Hardiness Zone: 11-Sep

Growth Habit: Vine

Origin: Eastern and southern Asia



FNAI

Description: Semi-woody vine, climbing by twining, or prostrate and rooting at the nodes, stems to 7 m or more long. Leaves opposite (rarely in whorls of 3), stipules conspicuous, petioles to 6 cm long, blades entire, oval to linear-lanceolate, 2-11 cm long, hairy or glabrous, often lobed at the base. Leaves and stems with fetid odor, especially when crushed. Flowers small, pedunculate, corolla tubular with 5 spreading lobes grayish pink or pale lilac with darker throat, outer surface pubescent. Fruit shiny, yellowish-orange, subglobose, to 7mm wide capsule. Seeds 2, black, round, not winged.

Note: Similar in appearance to *Paederia cruddasiana*.

Habitat: Sandhill, floodplain, upland mixed forest.

Florida Introduction Date: Before 1897.

Control Methods: .

Chemical: Foliar (3%-5% glyphosate), basal bark (10% trichlopyr ester). Within 2-4 weeks re-treat the area with basal applications of 10% trichlopyr ester. This second treatment can be time-consuming because many underground runners sprout. The area should continue to be monitored (IFAS)..

Useful Resources:

Langeland, K.A., J.A. Ferrell, B. Sellers, G.E. MacDonald, and R.K. Stocker. 2011. Integrated management of non-native plants in natural areas of Florida. EDIS publication SP 242. University of Florida, Gainesville, Florida.

Comments: Recently spreading into parts of North Florida.